arXiv:0805.2729v1 [math.AC] 18 May 2008

Intersections of base rings associated to transversal polymatroids

Alin Ştefan
"Petroleum and Gas" University of Ploieşti, Romania

Abstract

In this paper, we study when the intersection of base rings associated to some transversal polymatroids from [5] is the base ring of a transversal polymatroid. This intersection is a Gorenstein ring and we compute its a-invariant.

Key Words: Base ring, transversal polymatroid, equations of a cone, a-invariant, canonical module, Hilbert series.
2000 Mathematics Subject Classification:
Primary 13P10, Secondary 13H10, 13D40, 13A02.

1 Introduction

The discrete polymatroids and their base rings are studied recently in many papers (see [3], [4], [7], [8]). It is important to give conditions when the base ring associated to a transversal polymatroid is Gorenstein (see [3]). In [5] we introduced a class of such base rings. In this paper we note that an intersection of such base rings (introduced in [5]) is Gorenstein and give necessary and sufficient conditions for the intersection of two base rings from [5] to be still a base ring of a transversal polymatroid. Also, we compute the a-invariant of those base rings. The results presented were discovered by extensive computer algebra experiments performed with *Normaliz* [2].

2 Preliminaries

Let $n \in \mathbb{N}$, $n \geq 3$, $\sigma \in S_n$, $\sigma = (1, 2, \ldots, n)$ the cycle of length n, $[n] := \{1, 2, \ldots, n\}$, $\sigma^t[i] := \{\sigma^t(1), \ldots, \sigma^t(i)\}$ for any $1 \leq i \leq n-1$ and $\{e_i\}_{1 \leq i \leq n}$ be the canonical base of $\mathbb{R}^n$. For a vector $x \in \mathbb{R}^n$, $x = (x_1, \ldots, x_n)$, we will denote $\mid x \mid := x_1 + \ldots + x_n$. If x^a is a monomial in $K[x_1, \ldots, x_n]$ we set $\log(x^a) = a$. Given a set A of monomials, the *log set of* A, denoted $\log(A)$, consists of all $\log(x^a)$ with $x^a \in A$.
If A_i are some nonempty subsets of $[n]$ for $1 \leq i \leq m$, $m \geq 2$, $\mathcal{A} = \{A_1, \ldots, A_m\}$, then the set of the vectors $\sum_{k=1}^{m} e_{i_k}$ with $i_k \in A_k$ is the base of a polymatroid, called the *transversal polymatroid presented* by $\mathcal{A}$. The *base ring* of a transversal polymatroid presented by $\mathcal{A}$ is the ring

$$K[\mathcal{A}] := K[x_{i_1} \cdots x_{i_m} \mid i_j \in A_j, 1 \leq j \leq m].$$

We know by [3] that the $K-$ algebra $K[\mathcal{A}]$ is normal and hence Cohen-Macaulay. From [5] we know that the transversal polymatroid presented by

$$\mathcal{A} = \{A_k \mid A_{\sigma^t(k)} = [n],\ if\ k \in [i] \cup \{n\}\ and\ A_{\sigma^t(k)} = [n] \setminus \sigma^t[i],\ if\ k \in [n-1] \setminus [i]\}$$

has the base ring associated $K[\mathcal{A}]$ a Gorenstein ring for any $1 \leq i \leq n-1$ and $0 \leq t \leq n-1$.
We put

$$\nu_{\sigma^t[i]} = -(n-i-1) \sum_{k=1}^{i} e_{\sigma^t(k)} + (i+1) \sum_{k=i+1}^{n} e_{\sigma^t(k)}$$

for any $0 \leq t \leq n-1$ and $1 \leq i \leq n-1$.
The cone generated by the exponent vectors of the monomials defining the base ring $K[\mathcal{A}]$, $A := \{log(x_{j_1} \cdots x_{j_n}) \mid j_k \in A_k, k \in [n]\} \subset \mathbb{N}^n$, has the irreducible representation:

$$\mathbb{R}_+ A = \bigcap_{a \in N} H_a^+,$$

where $N = \{\nu_{\sigma^t[i]},\ \nu_{\sigma^k[n-1]} \mid 0 \leq k \leq n-1\}$.
It is easy to see that for any $1 \leq i \leq n-1$ and $0 \leq t \leq n-1$

$$A = \{\alpha \in \mathbb{N}^n \mid \ \mid \alpha \mid = n,\ 0 \leq \alpha_{t+1} + \ldots + \alpha_{t+i} \leq i+1\},\ if\ i+t \leq n$$

and

$$A = \{\alpha \in \mathbb{N}^n \mid \ \mid \alpha \mid = n,\ 0 \leq \sum_{s=1}^{i+t-n} \alpha_s + \sum_{s=t+1}^{n} \alpha_s \leq i+1\},\ if\ i+t\ >\ n.$$

We denote by $\{x_{i_1}, \ldots, x_{i_s}\}^r$ all monomials of degree r with the indeterminates $x_{i_1}, \ldots, x_{i_s}$.

3 Intersection of cones of dimension n with $n+1$ facets

Let $r \geq 2$, $1 \leq i_1, \ldots, i_r \leq n-2$, $0 = t_1 \leq t_2, \ldots, t_r \leq n-1$ and we consider r presentations of transversal polymatroids:

$$\mathcal{A}_s = \{A_{s,k} \mid A_{s,\sigma^{t_2}(k)} = [n],\ if\ k \in [i_2] \cup \{n\},\ A_{s,\sigma^{t_2}(k)} = [n] \setminus \sigma^{t_2}[i_2],\ if\ k \in [n-1] \setminus [i_2]\}$$

for any $1 \leq s \leq r$. From [5] we know that the base rings $K[\mathcal{A}_s]$ are Gorenstein rings and the cones generated by the exponent vectors of the monomials defining the base ring associated to a transversal polymatroid presented by $\mathcal{A}_s$ are :

$$\mathbb{R}_+ A_s = \bigcap_{a \in N_s} H_a^+,$$

where $N_s = \{\nu_{\sigma^{t_s}[i_s]},\ \nu_{\sigma^k[n-1]} \mid 0 \leq k \leq n-1\}$, $A_s = \{log(x_{j_1} \cdots x_{j_n}) \mid j_k \in A_{s,k}, 1 \leq k \leq n\} \subset \mathbb{N}^n$

for any $1 \leq s \leq r$.
We denote by $K[A_1 \cap \ldots \cap A_r]$, $K-algebra$ generated by x^α with $\alpha \in A_1 \cap \ldots \cap A_r$.
It is clear that the cone

$$\mathbb{R}_+(A_1 \cap \ldots \cap A_r) \subseteq \mathbb{R}_+ A_1 \cap \ldots \cap \mathbb{R}_+ A_r = \bigcap_{a \in N_1 \cup \ldots \cup N_r} H_a^+.$$

Conversely, since

$$A_1 \cap \ldots \cap A_r = \{\alpha \in \mathbb{N}^n \mid \ \mid \alpha \mid = n,\ H_{\nu_{\sigma^{t_s}[i_s]}}(\alpha) \geq 0,\ for\ any\ 1 \leq s \leq r\},$$

we have that

$$\mathbb{R}_+(A_1 \cap \ldots \cap A_r) \supseteq \bigcap_{a \in N_1 \cup \ldots \cup N_r} H_a^+$$

and so,

$$\mathbb{R}_+(A_1 \cap \ldots \cap A_r) = \bigcap_{a \in N_1 \cup \ldots \cup N_r} H_a^+.$$

We claim that the intersection

$$\bigcap_{a \in N_1 \cup \ldots \cup N_r} H_a^+$$

is the irreducible representation of the cone $\mathbb{R}_+(A_1 \cap \ldots \cap A_r)$.
We prove by induction on $r \geq 1$. If $r = 1$, then the intersection

$$\bigcap_{a \in N_1} H_a^+$$

is the irreducible representation of the cone $\mathbb{R}_+(A_1)$. (see [5], Lemma 4.1)
If $r > 1$ then we have two cases to study:
1) if we delete, for some $0 \leq k \leq n-1$, the hyperplane with the normal $\nu_{\sigma^k[n-1]}$, then a coordinate of a $\log(x_{j_1} \cdots x_{j_{i_1}} x_{j_{i_1+1}} \cdots x_{j_{n-1}} x_{j_n})$ would be negative, which is impossible;
2) if we delete, for some $1 \leq s \leq r$, the hyperplane with the normal $\nu_{\sigma^{t_s}[i_s]}$, then

$$\bigcap_{a \in N_1 \cup \ldots \cup N_{s-1} \cup N_{s+1} \ldots N_r} H_a^+$$

is by induction the irreducible representation of the cone $\mathbb{R}_+(A_1 \cap \ldots \cap A_{s-1} \cap A_{s+1} \ldots \cap A_r)$, which is diffrent from $\mathbb{R}_+(A_1 \cap \ldots \cap A_r)$. Hence, the intersection

$$\bigcap_{a \in N_1 \cup \ldots \cup N_r} H_a^+$$

is the irreducible representation of the cone $\mathbb{R}_+(A_1 \cap \ldots \cap A_r)$.

Lemma 3.1. *The $K-$ algebra $K[A_1 \cap \ldots \cap A_r]$ is a Gorenstein ring.*

Proof. We will show that the canonical module $\omega_{K[A_1 \cap \ldots \cap A_r]}$ is generated by $(x_1 \cdots x_n)K[A_1 \cap \ldots \cap A_r]$. Since the semigroups $\mathbb{N}(A_t)$ are normal for any $1 \leq t \leq r$, it follows that $\mathbb{N}(A_1 \cap \ldots \cap A_r)$ is normal. Then the $K-$ algebra $K[A_1 \cap \ldots \cap A_r]$ is normal (see [1] Theorem 6.1.4. p. 260) and using the *Danilov − Stanley* theorem we get that the canonical module $\omega_{K[A_1 \cap \ldots \cap A_r]}$ is

$$\omega_{K[A_1 \cap \ldots \cap A_r]} = (\{x^\alpha \mid \alpha \in \mathbb{N}(A_1 \cap \ldots \cap A_r) \cap ri(\mathbb{R}_+(A_1 \cap \ldots \cap A_r))\}).$$

Let d_t be the greatest common divisor of n *and* $i_t + 1$, $gcd(n, i_t + 1) = d_t$, for any $1 \leq t \leq r$.
For any $1 \leq s \leq r$, there exist two possibilities for the equation of the facet $H_{\nu_{\sigma^{t_s}[i_s]}}$:
1) If $i_s + t_s \leq n$, then the equation of the facet $H_{\nu_{\sigma^{t_s}[i_s]}}$ is:

$$H_{\nu_{\sigma^{t_s}[i_s]}}(y): \ \frac{(i_s+1)}{d_s}\sum_{k=1}^{t_s} y_k - \frac{(n-i_s-1)}{d_s}\sum_{k=t_s+1}^{t_s+i_s} y_k + \frac{(i_s+1)}{d_s}\sum_{k=t_s+i_s+1}^{n} y_k = 0.$$

2) If $i_s + t_s > n$, then the equation of the facet $H_{\nu_{\sigma^{t_s}[i_s]}}$ is:

$$H_{\nu_{\sigma^{t_s}[i_s]}}(y): \ -\frac{(n-i_s-1)}{d_s}\sum_{k=1}^{i_s+t_s-n} y_k + \frac{(i_s+1)}{d_s}\sum_{k=i_s+t_s-n+1}^{t_s} y_k - \frac{(n-i_s-1)}{d_s}\sum_{k=t_s+1}^{n} y_k = 0.$$

The relative interior of the cone $\mathbb{R}_+(A_1 \cap \ldots \cap A_r)$ is:

$$ri(\mathbb{R}_+(A_1 \cap \ldots \cap A_r)) = \{y \in \mathbb{R}^n \mid y_k > 0,\ H_{\nu_{\sigma^{t_s}[i_s]}}(y) > 0 \ for\ any\ 1 \leq k \leq n\ and\ 1 \leq s \leq r\}$$

We will show that

$$\mathbb{N}(A_1 \cap \ldots \cap A_r) \cap ri(\mathbb{R}_+(A_1 \cap \ldots \cap A_r)) = (1, \ldots, 1) + (\mathbb{N}(A_1 \cap \ldots \cap A_r) \cap \mathbb{R}_+(A_1 \cap \ldots \cap A_r)).$$

It is clear that $ri(\mathbb{R}_+(A_1 \cap \ldots \cap A_r)) \supset (1, \ldots, 1) + \mathbb{R}_+(A_1 \cap \ldots \cap A_r)$.
If $(\alpha_1, \alpha_2, \ldots, \alpha_n) \in \mathbb{N}(A_1 \cap \ldots \cap A_r) \cap ri(\mathbb{R}_+(A_1 \cap \ldots \cap A_r))$, then $\alpha_k \geq 1$ for any $1 \leq k \leq n$ and for any $1 \leq s \leq r$ we have

$$\frac{(i_s+1)}{d_s}\sum_{k=1}^{t_s} \alpha_k - \frac{(n-i_s-1)}{d_s}\sum_{k=t_s+1}^{t_s+i_s} \alpha_k + \frac{(i_s+1)}{d_s}\sum_{k=t_s+i_s+1}^{n} \alpha_k \geq 1,\ if\ i_s + t_s \leq n$$

or

$$-\frac{(n-i_s-1)}{d_s}\sum_{k=1}^{i_s+t_s-n}\alpha_k+\frac{(i_s+1)}{d_s}\sum_{k=i_s+t_s-n+1}^{t_s}\alpha_k-\frac{(n-i_s-1)}{d_s}\sum_{k=t_s+1}^{n}\alpha_k\geq 1,\ if\ i_s+t_s\ >\ n$$

and

$$\sum_{k=1}^{n}\alpha_k=t\ n\ for\ some\ t\geq 1.$$

We claim that there exist $(\beta_1,\beta_2,\ldots,\beta_n)\in\mathbb{N}(A_1\cap\ldots\cap A_r)\cap\mathbb{R}_+(A_1\cap\ldots\cap A_r)$ such that $(\alpha_1,\alpha_2,\ldots,\alpha_n)=(\beta_1+1,\beta_2+1,\ldots,\beta_n+1)$. Let $\beta_k=\alpha_k-1$ for all $1\leq k\leq n$.
It is clear that $\beta_k\geq 0$ and for any $1\leq s\leq r$,

$$H_{\nu_{\sigma^{t_s}[i_s]}}(\beta)=H_{\nu_{\sigma^{t_s}[i_s]}}(\alpha)-H_{\nu_{\sigma^{t_s}[i_s]}}(1,\ldots,1)=H_{\nu_{\sigma^{t_s}[i_s]}}(\alpha)-\frac{n}{d_s}.$$

If $H_{\nu_{\sigma^{t_s}[i_s]}}(\beta)=j_s,\ for\ some\ 1\leq s\leq r\ and\ 1\leq j_s\leq\frac{n}{d_s}-1$, then we will get a contadiction. Indeed, since n divides $\sum_{k=1}^{n}\alpha_k$, it follows $\frac{n}{d_s}$ divides j_s, which is false.
Hence, we have $(\beta_1,\beta_2,\ldots,\beta_n)\in\mathbb{N}(A_1\cap\ldots\cap A_r)\cap\mathbb{R}_+(A_1\cap\ldots\cap A_r)$ and $(\alpha_1,\alpha_2,\ldots,\alpha_n)\in\mathbb{N}(A_1\cap\ldots\cap A_r)\cap ri(\mathbb{R}_+(A_1\cap\ldots\cap A_r))$.
Since $\mathbb{N}(A_1\cap\ldots\cap A_r)\cap ri(\mathbb{R}_+(A_1\cap\ldots\cap A_r))=\ (1,\ldots,1)+(\mathbb{N}(A_1\cap\ldots\cap A_r)\cap\mathbb{R}_+(A_1\cap\ldots\cap A_r))$, we get that $\omega_{K[A_1\cap\ldots\cap A_r]}=\ (x_1\cdots x_n)K[A_1\cap\ldots\cap A_r]$. ☐

Let S be a standard graded $K-algebra$ over a field K. Recall that the $a-invariant$ of S, denoted $a(S)$, is the degree as a rational function of the Hilbert series of S, see for instance ([6], $p.$ 99). If S is $Cohen-Macaulay$ and ω_S is the canonical module of S, then

$$a(S)=\ -\ min\ \{i\mid(\omega_S)_i\neq 0\},$$

see ([1], p. 141) and ([6], Proposition 4.2.3). In our situation $S=K[A_1\cap\ldots\cap A_r]$ is normal and consequently $Cohen-Macaulay$, thus this formula applies. As consequence of $Lemma$ 3.1. we have the following:

Corollary 3.2. *The* $a-invariant$ *of* $K[A_1\cap\ldots\cap A_r]$ *is* $a(K[A_1\cap\ldots\cap A_r])=\ -1$.

Proof. Let $\{x^{\alpha_1},\ldots,x^{\alpha_q}\}$ be the generators of $K-algebra$ $K[A_1\cap\ldots\cap A_r]$. $K[A_1\cap\ldots\cap A_r]$ is standard graded algebra with the grading

$$K[A_1\cap\ldots\cap A_r]_i=\sum_{|c|=i}K(x^{\alpha_1})^{c_1}\cdots(x^{\alpha_q})^{c_q},\ where\ \mid c\mid=c_1+\ldots+c_q.$$

Since $\omega_{K[A_1\cap\ldots\cap A_r]}=\ (x_1\cdots x_n)K[A_1\cap\ldots\cap A_r]$ it follows that $min\ \{i\mid(\omega_{K[A_1\cap\ldots\cap A_r]})_i\neq 0\}=1$ and so, $a(K[A_1\cap\ldots\cap A_r])=\ -1$. ☐

4 When $K[A\cap B]$ is the base ring associated to some transversal polymatroid?

Let $n\geq 2$, and we consider two transversal polymatroids presented by: $\mathcal{A}=\{A_1,\ldots,A_n\}$ respectively $\mathcal{B}=\{B_1,\ldots,B_n\}$. Let A and B be the set of exponent vectors of monomials defining the base rings $K[\mathcal{A}]$, respectively $K[\mathcal{B}]$ and $K[A\cap B]$ the $K-algebra$ generated by x^α with $\alpha\in A\cap B$.
Question: There exists a transversal polymatroid such that its base ring is the $K-algebra$ $K[A\cap B]$?
In the following we will give two suggestive examples.
Example **1.** Let $n=4$, $\mathcal{A}=\{A_1,A_2,A_3,A_4\}$, $\mathcal{B}=\{B_1,B_2,B_3,B_4\}$, where $A_1=A_4=B_2=$

$B_3 = \{1,2,3,4\}$, $A_2 = A_3 = \{2,3,4\}$, $B_1 = B_4 = \{1,3,4\}$ and $K[\mathcal{A}]$, $K[\mathcal{B}]$ the base rings associated to transversal polymatroids presented by $\mathcal{A}$, respectively $\mathcal{B}$. It is easy to see that the generators set of $K[\mathcal{A}]$, respectively $K[\mathcal{B}]$ are given by $A = \{y \in \mathbb{N}^4 \mid \ \mid y \mid = 4,\ 0 \le y_1 \le 2,\ y_k \ge 0,\ 1 \le k \le 4\}$, respectively $B = \{y \in \mathbb{N}^4 \mid \ \mid y \mid = 4,\ 0 \le y_2 \le 2,\ y_k \ge 0,\ 1 \le k \le 4\}$. We show that the $K-$ algebra $K[A \cap B]$ is the base ring of the transversal polymatroid presented by $\mathcal{C} = \{C_1, C_2, C_3, C_4\}$, where $C_1 = C_4 = \{1,3,4\},\ C_2 = C_3 = \{2,3,4\}$.

Since the base ring associated to the transversal polymatroid presented by $\mathcal{C}$ has the exponent set $C = \{y \in \mathbb{N}^4 \mid \ \mid y \mid = 4,\ 0 \le y_1 \le 2,\ 0 \le y_2 \le 2,\ y_k \ge 0,\ 1 \le k \le 4\}$, it follows that $K[A \cap B] = K[\mathcal{C}]$. Thus, in this example $K[A \cap B]$ is the base ring of a transversal polymatroid.

***Example* 2.** Let $n = 4$, $\mathcal{A} = \{A_1, A_2, A_3, A_4\}$, $\mathcal{B} = \{B_1, B_2, B_3, B_4\}$ where $A_1 = A_2 = A_4 = B_1 = B_2 = B_3 = \{1,2,3,4\}$, $A_3 = \{3,4\}$, $B_4 = \{1,4\}$ and $K[\mathcal{A}]$, $K[\mathcal{B}]$ the base rings associated to the transversal polymatroids presented by $\mathcal{A}$, respectively $\mathcal{B}$. It is easy to see that the generators set of $K[\mathcal{A}]$, respectively $K[\mathcal{B}]$ are $A = \{y \in \mathbb{N}^4 \mid \ \mid y \mid = 4,\ 0 \le y_1 + y_2 \le 3,\ y_k \ge 0,\ 1 \le k \le 4\}$, respectively $B = \{y \in \mathbb{N}^4 \mid \ \mid y \mid = 4,\ 0 \le y_2 + y_3 \le 3,\ y_k \ge 0,\ 1 \le k \le 4\}$. We claim that there exists no transversal polymatroid $\mathcal{P}$ such that $K-$ algebra $K[A \cap B]$ is its base ring. Suppose, on the contrary, let $\mathcal{P}$ be presented by $\mathcal{C} = \{C_1, C_2, C_3, C_4\}$ with each $C_k \subset [4]$. Since $(3,0,1,0), (3,0,0,1) \in \mathcal{P}$ and $(3,1,0,0) \notin \mathcal{P}$, we may assume that changing the numerotation of $\{C_i\}_{i=\overline{1,4}}$ that $1 \in C_1, 1 \in C_2, 1 \in C_4$ and $C_3 = \{3,4\}$. Since $(0,3,0,1) \in \mathcal{P}$, we assume that $2 \in C_1, 2 \in C_2, 2 \in C_4$. Hence $(0,3,1,0) \in \mathcal{P}$, a contradiction.

Let $1 \le i_1, i_2 \le n-2$, $0 \le t_2 \le n-1$ and $\tau \in S_{n-2}$, $\tau = (1,2,\ldots,n-2)$ the cycle of length $n-2$. We consider two transversal polymatroids presented by:

$$\mathcal{A} = \{A_k \mid A_k = [n],\ if\ k \in [i_1] \cup \{n\},\ A_k = [n] \setminus [i_1],\ if\ k \in [n-1] \setminus [i_1]\}$$

and

$$\mathcal{B} = \{B_k \mid B_{\sigma^{t_2}(k)} = [n],\ if\ k \in [i_2] \cup \{n\},\ B_{\sigma^{t_2}(k)} = [n] \setminus \sigma^{t_2}[i_2],\ if\ k \in [n-1] \setminus [i_2]\}$$

such that A, respectively B is the exponent vectors of the monomials defining the base rings associated to transversal polymatroid presented by $\mathcal{A}$, respectively $\mathcal{B}$. From [5] we know that the base rings $K[\mathcal{A}]$, respectively $K[\mathcal{B}]$ are Gorenstein rings and the cones generated by the exponent vectors of the monomials defining the base ring associated to the transversal polymatroids presented by $\mathcal{A}$, respectively $\mathcal{B}$ are :

$$\mathbb{R}_+ A = \bigcap_{a \in N_1} H_a^+, \quad \mathbb{R}_+ B = \bigcap_{a \in N_2} H_a^+,$$

where $N_1 = \{\nu_{\sigma^0[i_1]},\ \nu_{\sigma^k[n-1]} \mid 0 \le k \le n-1\}$, $N_2 = \{\nu_{\sigma^{t_2}[i_2]},\ \nu_{\sigma^k[n-1]} \mid 0 \le k \le n-1\}$,

$A = \{log(x_{j_1} \cdots x_{j_n}) \mid j_k \in A_k, 1 \le k \le n\} \subset \mathbb{N}^n\ and\ B = \{log(x_{j_1} \cdots x_{j_n}) \mid j_k \in B_k, 1 \le k \le n\} \subset \mathbb{N}^n$.

It is easy to see that $A = \{\alpha \in \mathbb{N}^n \mid \ 0 \le \alpha_1 + \ldots + \alpha_{i_1} \le i_1 + 1\ and\ \mid \alpha \mid = n\}$ and
$B = \{\alpha \in \mathbb{N}^n \mid \ 0 \le \alpha_{t_2+1} + \ldots + \alpha_{t_2+i_2} \le i_2 + 1\ and\ \mid \alpha \mid = n\},\ if\ i_2 + t_2 \le n$
or

$$B = \{\alpha \in \mathbb{N}^n \mid 0 \le \sum_{s=1}^{i_2+t_2-n} \alpha_s + \sum_{s=t_2+1}^{n} \alpha_s \le i_2 + 1\ and\ \mid \alpha \mid = n\},\ if\ i_2 + t_2 \ge n.$$

For any base ring $K[\mathcal{A}]$ of a transversal polymatroid presented by $\mathcal{A} = \{A_1, \ldots, A_n\}$ we associate a $(n \times n)$ square tiled by closed unit subsquares, called **boxes**, colored with colors, "*white*" and "*black*", as follows: the box of coordinate (i,j) is "*white*" if $j \in A_i$, otherwise the box is "*black*". We will call this square the **polymatroidal diagram** associated to the presentation $\mathcal{A} = \{A_1, \ldots, A_n\}$.

Next we give necesary and sufficient conditions such that the $K-algebra$ $K[A \cap B]$ is the base ring associated to some transversal polymatroid.

Theorem 4.1. *Let* $1 \leq i_1, i_2 \leq n-2$, $0 \leq t_2 \leq n-1$. *We consider two presentation of transversal poymatroids presented by:* $\mathcal{A} = \{A_k \mid A_k = [n],\ if\ k \in [i_1] \cup \{n\},\ A_k = [n] \setminus [i_1],\ if\ k \in [n-1] \setminus [i_1]\}$ *and* $\mathcal{B} = \{B_k \mid B_{\sigma^{t_2}(k)} = [n],\ if\ k \in [i_2] \cup \{n\},\ B_{\sigma^{t_2}(k)} = [n] \setminus \sigma^{t_2}[i_2],\ if\ k \in [n-1] \setminus [i_2]\}$ *such that* A, *respectively* B *are the exponent vectors of the monomials defining the base ring associated to the transversal polymatroids presented by* $\mathcal{A}$, *respectively* $\mathcal{B}$.
Then, the $K-algebra$ $K[A \cap B]$ *is the base ring associated to a transversal polymatroid if and only if one of the following conditions hold:*
a) $i_1 = 1$.
b) $i_1 \geq 2$ *and* $t_2 = 0$.
c) $i_1 \geq 2$ *and* $t_2 = i_1$.
d) $i_1 \geq 2$, $1 \leq t_2 \leq i_1 - 1$ *and* $i_2 \in \{1, \ldots, i_1 - t_2\} \cup \{n - t_2, \ldots, n-2\}$;
e) $i_1 \geq 2$, $i_1 + 1 \leq t_2 \leq n-1$ *and* $i_2 \in \{1, \ldots, n - t_2\} \cup \{n - t_2 + i_1, \ldots, n-2\}$.

The proof follows from the following three lemmas.

Lemma 4.2. *Let* A *and* B *be like above. If* $i_1 \geq 2$, $1 \leq t_2 \leq i_1 - 1$, *then the* $K-algebra$ $K[A \cap B]$ *is the base ring associated to some transversal polymatroid if and only if* $i_2 \in \{1, \ldots, i_1 - t_2\} \cup \{n - t_2, \ldots, n-2\}$.

Proof. " $\Leftarrow$ " Let $i_2 \in \{1, \ldots, i_1 - t_2\} \cup \{n - t_2, \ldots, n-2\}$. We will prove that there exists a transversal polymatroid $\mathcal{P}$ presented by $\mathcal{C} = \{C_1, \ldots, C_n\}$ such that the base ring associated to $\mathcal{P}$ is $K[A \cap B]$.
We have two cases to study:
Case 1. If $i_2 + t_2 \leq i_1$, then let $\mathcal{P}$ be the transversal polymatroid presented by $\mathcal{C} = \{C_1, \ldots, C_n\}$, where

$$C_1 = \ldots = C_{i_2} = C_n = [n],$$

$$C_{i_2+1} = \ldots = C_{i_1} = [n] \setminus \sigma^{t_2}[i_2],$$

$$C_{i_1+1} = \ldots = C_{n-1} = [n] \setminus [i_1].$$

The polymatroidal diagram associated is the following:



It is easy to see that the base ring associated to the transversal polymatroid $\mathcal{P}$ presented by $\mathcal{C}$ is generated by the following set of monomials:

$$\{x_{t_2+1}, \ldots, x_{t_2+i_2}\}^{i_2+1-k}\{x_1, \ldots, x_{t_2}, x_{t_2+i_2+1}, \ldots, x_{i_1}\}^{i_1-i_2+k-s}\{x_{i_1+1}, \ldots, x_n\}^{n-1-i_1+s}$$

for any $0 \leq k \leq i_2 + 1$ and $0 \leq s \leq i_1 - i_2 + k$. If $x^\alpha \in K[\mathcal{C}]$, $\alpha = (\alpha_1, \ldots, \alpha_n) \in \mathbb{N}^n$, then there exists $0 \leq k \leq i_2 + 1$ and $0 \leq s \leq i_1 - i_2 + k$ such that

$$\alpha_{t_2+1} + \ldots + \alpha_{t_2+i_2} = i_2 + 1 - k \ and \ \alpha_1 + \ldots + \alpha_{i_1} = i_1 + 1 - s$$

and thus, $K[\mathcal{C}] \subset K[A \cap B]$.
Conversely, if $\alpha \in A \cap B$ then $\alpha_{t_2+1} + \ldots + \alpha_{t_2+i_2} \leq i_2 + 1$ *and* $\alpha_1 + \ldots + \alpha_{i_1} \leq i_1 + 1$ and thus there exists $0 \leq k \leq i_2 + 1$ and $0 \leq s \leq i_1 - i_2 + k$ such that

$$x^\alpha \in \{x_{t_2+1}, \ldots, x_{t_2+i_2}\}^{i_2+1-k}\{x_1, \ldots, x_{t_2}, x_{t_2+i_2+1}, \ldots, x_{i_1}\}^{i_1-i_2+k-s}\{x_{i_1+1}, \ldots, x_n\}^{n-1-i_1+s}.$$

Thus, $K[\mathcal{C}] \supset K[A \cap B]$ and so $K[\mathcal{C}] = K[A \cap B]$.
Case 2. If $i_2 + t_2 > i_1$, then it follows that $i_2 \geq n - t_2$ and $n - i_2 \leq t_2 < i_1$. Let $\mathcal{P}$ be the transversal polymatroid presented by $\mathcal{C} = \{C_1, \ldots, C_n\}$, where

$$C_1 = \ldots = C_{n-i_2-1} = [n] \setminus \sigma^{t_2}[i_2],$$

$$C_{n-i_2} = \ldots = C_{i_1} = C_n = [n],$$

$$C_{i_1+1} = \ldots = C_{n-1} = [n] \setminus [i_1].$$

The polymatroidal diagram associated is the following:



It is easy to see that the base ring associated to the transversal polymatroid $\mathcal{P}$ presented by $\mathcal{C}$ is generated by the following set of monomials:

$$\{x_{i_2+t_2-n+1}, \ldots, x_{t_2}\}^{i_1+1-k}\{x_1, \ldots, x_{i_2+t_2-n}, x_{t_2+1}, \ldots, x_{i_1}\}^{k-s}\{x_{i_1+1}, \ldots, x_n\}^{n-1-i_1+s}$$

for any $0 \leq k \leq i_1 + i_2 - n + 2$ and $0 \leq s \leq k$. Since $i_2 + t_2 \geq n$ and $0 \leq s \leq k \leq i_1 + i_2 - n + 2$ it follows that for any $x^\alpha \in K[\mathcal{C}]$ we have $\alpha_1 + \ldots + \alpha_{i_1} = i_1 + 1 - s \leq i_1 + 1$, $\alpha_1 + \ldots + \alpha_{i_2+t_2-n} + \alpha_{t_2+1} + \ldots + \alpha_n = n - 1 - i_1 + k \leq i_2 + 1$ and thus, $K[\mathcal{C}] \subset K[A \cap B]$.
Conversely, if $\alpha \in A \cap B$ then $\alpha_1 + \ldots + \alpha_{i_1} \leq i_1 + 1$, $\alpha_1 + \ldots + \alpha_{i_2+t_2-n} + \alpha_{t_2+1} + \ldots + \alpha_n \leq i_2 + 1$ and thus there exists $0 \leq k \leq i_1 + i_2 - n + 2$ and $0 \leq s \leq k$ such that

$$x^\alpha \in \{x_{i_2+t_2-n+1}, \ldots, x_{t_2}\}^{i_1+1-k}\{x_1, \ldots, x_{i_2+t_2-n}, x_{t_2+1}, \ldots, x_{i_1}\}^{k-s}\{x_{i_1+1}, \ldots, x_n\}^{n-1-i_1+s}.$$

Thus, $K[\mathcal{C}] \supset K[A \cap B]$ and so $K[\mathcal{C}] = K[A \cap B]$.

" $\Rightarrow$ " Now suppose that there exists a transversal polymatroid $\mathcal{P}$ given by $\mathcal{C} = \{C_1, \ldots, C_n\}$ such that its associated base ring is $K[A \cap B]$ and we will prove that $i_2 \in \{1, \ldots, i_1 - t_2\} \cup \{n - t_2, \ldots, n - 2\}$.
Suppose, on the contrary, let $i_1 + 1 - t_2 \leq i_2 \leq n - t_2 - 1$. We have two cases to study:
Case $1'$. If $n - i_1 - 1 \leq i_2 + 1$, then since $(i_1 + 1)e_1 + (n - i_1 - 1)e_k \in \mathcal{P}$ for any $i_1 + 1 \leq k \leq n$ and $(i_1 + 1)e_1 + e_s + (n - i_1 - 2)e_k \notin \mathcal{P}$ for any $2 \leq s \leq i_1$ and $i_1 + 1 \leq k \leq n$, we may assume $1 \in C_1, \ldots, 1 \in C_{i_1}, 1 \in C_n$ and $C_{i_1+1} = \ldots = C_{n-1} = [n] \setminus [i_1]$. If $i_1 \leq i_2$, then since $(i_1 + 1)e_{t_2+1} + (n - i_1 - 1)e_{t_2+i_2+1} \in \mathcal{P}$, we may assume $t_2 + 1 \in C_1, \ldots, t_2 + 1 \in C_{i_1}, t_2 + 1 \in C_n$. Then $(i_1 + 1)e_{t_2+1} + (n - i_1 - 1)e_{i_1+1} \in \mathcal{P}$, which is false. If $i_1 > i_2$, then since $(i_2 + 1)e_{t_2+1} +$

$(n-i_1-1)e_{t_2+i_2+1} \in \mathcal{P}$, we may assume $t_2+1 \in C_1, \ldots, t_2+1 \in C_{i_2}, t_2+1 \in C_n$. Then $(i_2+1)e_{t_2+1}+(i_1-i_2)e_1+(n-i_1-1)e_{i_1+1} \in \mathcal{P}$, which is false.

Case $2'$. If $n-i_1-1 > i_2+1$, then since $(i_1+1)e_1+(i_2+1)e_{i_1+1}+(n-i_1-i_2-2)e_k \in \mathcal{P}$ for any $t_2+i_2+1 \leq k \leq n$ and $(i_1+1)e_1+e_s+(i_2+1)e_{i_1+1}+(n-i_1-i_2-3)e_k \notin \mathcal{P}$ for any $1 \leq s \leq i_1$ and $t_2+i_2+1 \leq k \leq n$, we may assume $1 \in C_1, \ldots, 1 \in C_{i_1}, 1 \in C_n$, $C_{i_1+1} = \ldots = C_{i_1+i_2+1} = [n] \setminus [i_1]$ and $C_{i_1+i_2+2} = \ldots = C_{n-1} = [n] \setminus [t_2+i_2]$. If $i_1 \leq i_2$, then since $(i_1+1)e_{t_2+1}+(n-i_1-1)e_{t_2+i_2+1} \in \mathcal{P}$, we may assume $t_2+1 \in C_1, \ldots, t_2+1 \in C_{i_1}, t_2+1 \in C_n$. Then $(i_1+1)e_{t_2+1}+(i_2+1)e_{i_1+1}+(n-i_1-i_2-2)e_{t_2+i_2+1} \in \mathcal{P}$, which is false. If $i_1 > i_2$, then since $(i_2+1)e_{t_2+1}+(i_1-i_2)e_1+(n-i_1-1)e_{t_2+i_2+1} \in \mathcal{P}$, we may assume $t_2+1 \in C_1, \ldots, t_2+1 \in C_{i_2}, t_2+1 \in C_n$. Then $(i_1-i_2)e_1+(i_2+1)e_{t_2+1}+(i_2+1)e_{t_2+i_2}+(n-i_1-i_2-2)e_{t_2+i_2+1} \in \mathcal{P}$, which is false. $\square$

Lemma 4.3. *Let A and B be like above. If $i_1 \geq 2$, $i_1+1 \leq t_2 \leq n-1$, then the $K-algebra$ $K[A \cap B]$ is the base ring associated to some transversal polymatroid if and only if $i_2 \in \{1, \ldots, n-t_2\} \cup \{n-t_2+i_1, \ldots, n-2\}$.*

Proof. " $\Leftarrow$ " Let $i_2 \in \{1, \ldots, n-t_2\} \cup \{n-t_2+i_1, \ldots, n-2\}$. We will prove that there exists a transversal polymatroid $\mathcal{P}$ presented by $\mathcal{C} = \{C_1, \ldots, C_n\}$ such that its associated base ring is $K[A \cap B]$. We distinct three cases to study:

Case 1. If $i_2+t_2 \leq n$ and $i_1+1+i_2 \neq n$, then let $\mathcal{P}$ be the transversal polymatroid presented by $\mathcal{C} = \{C_1, \ldots, C_n\}$, where

$$C_1 = \ldots = C_{i_1} = C_n = [n] \setminus \sigma^{t_2}[i_2],$$
$$C_{i_1+1} = \ldots = C_{i_1+i_2+1} = [n] \setminus [i_1],$$
$$C_{i_1+i_2+2} = \ldots = C_{n-1} = [n] \setminus ([i_1] \cup \sigma^{t_2}[i_2]).$$

The polymatroidal diagram associated is the following:



It is easy to see that the base ring $K[\mathcal{C}]$ associated to the transversal polymatroid $\mathcal{P}$ presented by $\mathcal{C}$ is generated by the following set of monomials:

$$\{x_1, \ldots, x_{i_1}\}^{i_1+1-k}\{x_{t_2+1}, \ldots, x_{t_2+i_2}\}^{i_2+1-s}\{x_{i_1+1}, \ldots, x_{t_2}, x_{t_2+i_2+1}, \ldots, x_n\}^{n-i_1-i_2-2+k+s}$$

for any $0 \leq k \leq i_1+1$ and $0 \leq s \leq i_2+1$. If $x^\alpha \in K[\mathcal{C}]$, $\alpha = (\alpha_1, \ldots, \alpha_n) \in \mathbb{N}^n$, then there exists $0 \leq k \leq i_1+1$ and $0 \leq s \leq i_2+1$ such that

$$\alpha_{t_2+1}+\ldots+\alpha_{t_2+i_2} = i_2+1-s \ , \quad \alpha_1+\ldots+\alpha_{i_1} = i_1+1-k$$

and thus, $K[\mathcal{C}] \subset K[A \cap B]$. Conversely, if $\alpha \in A \cap B$ then $\alpha_{t_2+1}+\ldots+\alpha_{t_2+i_2} \leq i_2+1$, $\alpha_1+\ldots+\alpha_{i_1} \leq i_1+1$; thus there exists $0 \leq k \leq i_1+1$ and $0 \leq s \leq i_2+1$ such that

$$\alpha_{t_2+1}+\ldots+\alpha_{t_2+i_2} = i_2+1-s \ , \quad \alpha_1+\ldots+\alpha_{i_1} = i_1+1-k$$

and since $\mid \alpha \mid = n$ it follows that

$$x^{\alpha} \in \{x_1, \ldots, x_{i_1}\}^{i_1+1-k}\{x_{t_2+1}, \ldots, x_{t_2+i_2}\}^{i_2+1-s}\{x_{i_1+1}, \ldots, x_{t_2}, x_{t_2+i_2+1}, \ldots, x_n\}^{n-i_1-i_2-2+k+s}.$$

Thus, $K[\mathcal{C}] \supset K[A \cap B]$ and so $K[\mathcal{C}] = K[A \cap B]$.

Case 2. If $i_2 + t_2 \leq n$ and $i_1 + 1 + i_2 = n$, then $t_2 = i_1 + 1$ and let $\mathcal{P}$ be the transversal polymatroid presented by $\mathcal{C} = \{C_1, \ldots, C_n\}$, where

$$C_1 = \ldots = C_{i_1} = [n] \setminus \sigma^{t_2}[i_2],$$

$$C_{i_1+1} = \ldots = C_{n-1} = [n] \setminus [i_1],$$

$$C_n = [n].$$

The polymatroidal diagram associated is the following:



It is easy to see that the base ring $K[\mathcal{C}]$ associated to the transversal polymatroid $\mathcal{P}$ presented by $\mathcal{C}$ is generated by the following set of monomials:

$$\{x_1, \ldots, x_{i_1}\}^{i_1+1-k}x_{i_1+1}^{n-i_1-1+k-s}\{x_{i_1+2}, \ldots, x_n\}^{s}$$

for any $0 \leq k \leq i_1 + 1$ and $0 \leq s \leq n - i_1$. If $x^{\alpha} \in K[\mathcal{C}]$, $\alpha = (\alpha_1, \ldots, \alpha_n) \in \mathbb{N}^n$, then there exists $0 \leq k \leq i_1 + 1$ and $0 \leq s \leq i_2 + 1 (= n - i_1)$ such that

$$\alpha_{t_2+1} + \ldots + \alpha_{t_2+i_2} = \alpha_{i_1+2} + \ldots + x_n = s \leq i_2 + 1 \ , \quad \alpha_1 + \ldots + \alpha_{i_1} = i_1 + 1 - k$$

and thus, $K[\mathcal{C}] \subset K[A \cap B]$. Conversely, if $\alpha \in A \cap B$ then $\alpha_{t_2+1} + \ldots + \alpha_{t_2+i_2} = \alpha_{i_1+2} + \ldots + x_n \leq i_2 + 1$ *and* $\alpha_1 + \ldots + \alpha_{i_1} \leq i_1 + 1$; thus there exists $0 \leq k \leq i_1 + 1$ and $0 \leq s \leq i_2 + 1$ such that

$$x^{\alpha} \in \{x_1, \ldots, x_{i_1}\}^{i_1+1-k}x_{i_1+1}^{n-i_1-1+k-s}\{x_{i_1+2}, \ldots, x_n\}^{s}.$$

Thus, $K[\mathcal{C}] \supset K[A \cap B]$ and so $K[\mathcal{C}] = K[A \cap B]$.

Case 3. If $i_2 + t_2 > n$, then let $\mathcal{P}$ be the transversal polymatroid presented by $\mathcal{C} = \{C_1, \ldots, C_n\}$, where

$$C_1 = \ldots = C_{i_1} = C_n = [n],$$

$$C_{i_1+1} = \ldots = C_{i_1+n-i_2-1} = [n] \setminus \sigma^{t_2}[i_2],$$

$$C_{i_1+n-i_2} = \ldots = C_{n-1} = [n] \setminus [i_1].$$

The polymatroidal diagram associated is the following:



Since $i_2+t_2>n$ it follows that $i_2+t_2\geq n+i_1$ and so $i_2-i_1\geq n-t_2\geq 1$. It is easy to see that the base ring $K[\mathcal{C}]$ associated to the transversal polymatroid $\mathcal{P}$ presented by $\mathcal{C}$ is generated by the following set of monomials:

$$\{x_1,\ldots,x_{i_1}\}^{i_1+1-k}\{x_{i_1+1},\ldots,x_{i_2+t_2-n},x_{t_2+1},\ldots,x_n\}^{i_2-i_1+k-s}\{x_{i_2+t_2-n+1},\ldots,x_{t_2}\}^{n-i_2-1+s}$$

for any $0\leq k\leq i_1+1$ and $0\leq s\leq i_2-i_1+k$. If $x^\alpha\in K[\mathcal{C}]$, $\alpha=(\alpha_1,\ldots,\alpha_n)\in\mathbb{N}^n$, then there exists $0\leq k\leq i_1+1$ and $0\leq s\leq i_2-i_1+k$ such that

$$\alpha_1+\ldots+\alpha_{i_1}=i_1+1-k\ ,\quad \alpha_1+\ldots+\alpha_{i_2+t_2-n}+\alpha_{t_2+1}+\ldots+\alpha_n=i_2+1-s$$

and thus, $K[\mathcal{C}]\subset K[A\cap B]$. Conversely, if $\alpha\in A\cap B$ then $\alpha_1+\ldots+\alpha_{i_1}\leq i_1+1$ *and* $\alpha_1+\ldots+\alpha_{i_2+t_2-n}+\alpha_{t_2+1}+\ldots+\alpha_n\leq i_2+1$. Then there exists $0\leq k\leq i_1+1$ and $0\leq s\leq i_2-i_1+k$ such that

$$\alpha_1+\ldots+\alpha_{i_1}=i_1+1-k\ ,\quad \alpha_1+\ldots+\alpha_{i_2+t_2-n}+\alpha_{t_2+1}+\ldots+\alpha_n=i_2+1-s$$

and since $\mid\alpha\mid=n$ it follows that

$$x^\alpha\in\{x_1,\ldots,x_{i_1}\}^{i_1+1-k}\{x_{i_1+1},\ldots,x_{i_2+t_2-n},x_{t_2+1},\ldots,x_n\}^{i_2-i_1+k-s}\{x_{i_2+t_2-n+1},\ldots,x_{t_2}\}^{n-i_2-1+s}.$$

Thus, $K[\mathcal{C}]\supset K[A\cap B]$ and so $K[\mathcal{C}]=K[A\cap B]$.

" $\Rightarrow$ " Now suppose that there exists a transversal polymatroid $\mathcal{P}$ presented by $\mathcal{C}=\{C_1,\ldots,C_n\}$ such that its associated base ring is $K[A\cap B]$ and we will prove that $i_2\in\{1,\ldots,n-t_2\}\cup\{n-t_2+i_1,\ldots,n-2\}$. Suppose, on the contrary, let $i_2\in\{n-t_2+1,\ldots,n-t_2+i_1-1\}$. We may assume that $1\notin C_{i_2+t_2-n+1},\ldots,1\notin C_{i_1},1\notin C_{i_1+1},\ldots,1\notin C_{n-1}$. If $i_1<i_2$, then $x_1^{i_1+1}x_{i_1+1}^{n-i_1-1}\in K[A\cap B]$. But $x_1^{i_1+1}x_{i_1+1}^{n-i_1-1}\notin K[\mathcal{C}]$ because the maximal power of x_1 in a minimal generator of $K[C]$ is $\leq i_2+t_2-n+1\leq i_1$, which is false. If $i_1\geq i_2$, then $x_1^{i_2+1}x_{i_1+1}^{n-i_2-1}\in K[A\cap B]$. But $x_1^{i_2+1}x_{i_1+1}^{n-i_2-1}\notin K[\mathcal{C}]$ because the maximal power of x_1 in a minimal generator of $K[C]$ is $\leq i_2+t_2-n+1\leq i_2$, which is false. Thus, $i_2\in\{1,\ldots,n-t_2\}\cup\{n-t_2+i_1,\ldots,n-2\}$. □

Lemma 4.4. *Let A and B like above. If either $i_1=1$ and $0\leq t_2\leq n-1$, or $i_1\geq 2$ and $t_2=i_1$, or $i_1\geq 2$ and $t_2=0$, then the $K-$algebra $K[A\cap B]$ is the base ring associated to some transversal polymatroid.*

Proof. We have three cases to study:
Case 1. $i_1=1$ and $0\leq t_2\leq n-1$. Then we distinct five subcases:
Subcase 1.*a*. If $t_2=0$, then we find a transversal polymatroid $\mathcal{P}$ like in the Subcase 3.*b*. when $i_1=1$.

Subcase 1.*b*. If $t_2 > 0$ *and* $t_2 + i_2 \leq n$ *with* $i_2 \neq n-2,\ n-3$, then let $\mathcal{P}$ be the transversal polymatroid presented by $\mathcal{C} = \{C_1, \ldots, C_n\}$, where

$$C_1 = C_n = [n] \setminus \sigma^{t_2}[i_2],$$

$$C_2 = \ldots = C_{i_2+2} = [n] \setminus [1],$$

$$C_{i_2+3} = \ldots = C_{n-1} = [n] \setminus \{\{1\} \cup \sigma^{t_2}[i_2]\}.$$

It is easy to see that the polymatroid $\mathcal{P}$ is the same like in *Lemma* 4.3 when $i_2 + t_2 \leq n$ *and* $i_1 + 1 + i_2 \neq n$. Thus $K[A \cap B] = K[\mathcal{C}]$.

Subcase 1.*c*. If $t_2 > 0$ *and* $t_2 + i_2 \leq n$ *with* $i_2 = n-2$, then let $\mathcal{P}$ be the transversal polymatroid presented by $\mathcal{C} = \{C_1, \ldots, C_n\}$, where

$$C_1 = [n] \setminus \sigma^{t_2}[n-2]\ ,\ C_n = [n],$$

$$C_2 = \ldots = C_{n-1} = [n] \setminus [1].$$

It is easy to see that the polymatroid $\mathcal{P}$ is the same like in *Lemma* 4.3. when $i_2 + t_2 \leq n$ *and* $i_1 + 1 + i_2 = n$. Thus $K[A \cap B] = K[\mathcal{C}]$.

Subcase 1.*d*. If $t_2 > 0$ *and* $t_2 + i_2 \leq n$ *with* $i_2 = n-3$, then let $\mathcal{P}$ be the transversal polymatroid presented by $\mathcal{C} = \{C_1, \ldots, C_n\}$, where

$$C_1 = C_n = [n] \setminus \sigma^{t_2}[n-3],$$

$$C_2 = \ldots = C_{n-1} = [n] \setminus [1], \qquad .$$

It is easy to see that the polymatroid $\mathcal{P}$ is the same like in *Lemma* 4.3. when $i_2 + t_2 \leq n$ *and* $i_1 + 1 + i_2 \neq n$. Thus $K[A \cap B] = K[\mathcal{C}]$.

Subcase 1.*e*. If $t_2 > 0$ and $t_2 + i_2 > n$, then let $\mathcal{P}$ be the transversal polymatroid presented by $\mathcal{C} = \{C_1, \ldots, C_n\}$, where

$$C_1 = C_n = [n],$$

$$C_2 = \ldots = C_{n-i_2} = [n] \setminus \sigma^{t_2}[i_2],$$

$$C_{n-i_2+1} = \ldots = C_{n-1} = [n] \setminus \{1\}.$$

It is easy to see that the polymatroid $\mathcal{P}$ is the same like in *Lemma* 4.3 when $i_2 + t_2 > n$ *and* $i_1 = 1$. Thus $K[A \cap B] = K[\mathcal{C}]$.

Case 2. $i_1 \geq 2$ and $t_2 = i_1$. Then we distinct three subcases:

Subcase 2.*a*. If $i_2 + t_2 < n-1$, then let $\mathcal{P}$ be the transversal polymatroid presented by $\mathcal{C} = \{C_1, \ldots, C_n\}$, where

$$C_1 = \ldots = C_{i_1} = C_n = [n] \setminus \sigma^{t_2}[i_2],$$

$$C_{i_1+1} = \ldots = C_{i_1+i_2+1} = [n] \setminus [i_1],$$

$$C_{i_1+i_2+2} = \ldots = C_{n-1} = [n] \setminus [i_1 + i_2].$$

It is easy to see that the polymatroid $\mathcal{P}$ is the same like in *Lemma* 4.3 when $i_2 + t_2 \leq n$ *and* $i_1 + 1 + i_2 \neq n$. Thus $K[A \cap B] = K[\mathcal{C}]$.

Subcase 2.*b*. If $i_2 + t_2 = n-1$, then let $\mathcal{P}$ be the transversal polymatroid presented by $\mathcal{C} = \{C_1, \ldots, C_n\}$, where

$$C_1 = \ldots = C_{i_1} = [n] \setminus \sigma^{t_2}[i_2],$$

$$C_{i_1+1} = \ldots = C_{n-1} = [n] \setminus [i_1],$$

$$C_n = [n].$$

It is easy to see that the polymatroid $\mathcal{P}$ is the same like in *Lemma* 4.3 when $i_2 + t_2 \leq n$ *and* $i_1 + 1 + i_2 = n$. Thus $K[A \cap B] = K[\mathcal{C}]$.

Subcase 2.*c*. If $i_2 + t_2 \geq n$, then let $\mathcal{P}$ be the transversal polymatroid presented by $\mathcal{C} = \{C_1, \ldots, C_n\}$, where

$$C_1 = \ldots = C_{n-i_2-1} = [n] \setminus \sigma^{t_2}[i_2],$$

$$C_{n-i_2} = \ldots = C_{i_1} = C_n = [n],$$

$$C_{i_1+1} = \ldots = C_{n-1} = [n] \setminus [i_1].$$

It is easy to see that the polymatroid $\mathcal{P}$ is the same like in *Lemma* 4.2 when $i_2 + t_2 > i_1$. Thus $K[A \cap B] = K[\mathcal{C}]$.

Case 3. $i_1 \geq 2$ and $t_2 = 0$. Then we distinct two subcases:

Subcase 3.*a*. If $i_2 \leq i_1$, then let $\mathcal{P}$ be the transversal polymatroid presented by $\mathcal{C} = \{C_1, \ldots, C_n\}$, where

$$C_1 = \ldots = C_{i_2} = C_n = [n],$$

$$C_{i_2+1} = \ldots = C_{i_1} = [n] \setminus [i_2],$$

$$C_{i_1+1} = \ldots = C_{n-1} = [n] \setminus [i_1].$$

It is easy to see that the polymatroid $\mathcal{P}$ is the same like in *Lemma* 4.2 when $i_2 + t_2 \leq i_1$. Thus $K[A \cap B] = K[\mathcal{C}]$.

Subcase 3.*b*. If $i_2 > i_1$, then let $\mathcal{P}$ be the transversal polymatroid presented by $\mathcal{C} = \{C_1, \ldots, C_n\}$, where

$$C_1 = \ldots = C_{i_1} = C_n = [n],$$

$$C_{i_1+1} = \ldots = C_{i_2} = [n] \setminus [i_1],$$

$$C_{i_2+1} = \ldots = C_{n-1} = [n] \setminus [i_2].$$

The polymatroidal diagram associated is the following:



It is easy to see that the base ring $K[\mathcal{C}]$ associated to the transversal polymatroid $\mathcal{P}$ presented by $\mathcal{C}$ is generated by the following set of monomials:

$$\{x_1, \ldots, x_{i_1}\}^{i_1+1-k}\{x_{i_1+1}, \ldots, x_{i_2}\}^{i_2-i_1+k-s}\{x_{i_2+1}, \ldots, x_n\}^{n-i_2+s-1}$$

for any $0 \leq k \leq i_1 + 1$ and $0 \leq s \leq i_2 - i_1 + k$. If $x^\alpha \in K[\mathcal{C}]$, $\alpha = (\alpha_1, \ldots, \alpha_n) \in \mathbb{N}^n$, then there exists $0 \leq k \leq i_1 + 1$ and $0 \leq s \leq i_2 - i_1 + k$ such that

$$\alpha_1 + \ldots + \alpha_{i_2} = i_2 + 1 - s \ and \ \alpha_1 + \ldots + \alpha_{i_1} = i_1 + 1 - k$$

and thus, $K[\mathcal{C}] \subset K[A \cap B]$. Conversely, if $\alpha \in A \cap B$ then $\alpha_1 + \ldots + \alpha_{i_2} \leq i_2 + 1$ *and* $\alpha_1 + \ldots + \alpha_{i_1} \leq i_1 + 1$ and so there exists $0 \leq k \leq i_1 + 1$ and $0 \leq s \leq i_2 - i_1 + k$ such that

$$\alpha_1 + \ldots + \alpha_{i_2} = i_2 + 1 - s \text{ and } \alpha_1 + \ldots + \alpha_{i_1} = i_1 + 1 - k$$

and since $\mid \alpha \mid = n$ it follows that

$$\{x_1, \ldots, x_{i_1}\}^{i_1+1-k}\{x_{i_1+1}, \ldots, x_{i_2}\}^{i_2-i_1+k-s}\{x_{i_2+1}, \ldots, x_n\}^{n-i_2+s-1}$$

Thus, $K[\mathcal{C}] \supset K[A \cap B]$ and so $K[\mathcal{C}] = K[A \cap B]$.

□

Acknowledgments: The author whould like to thank Professor Dorin Popescu for valuable suggestions and comments during the preparation of this paper.

References

[1] W. Bruns, J. Herzog, *Cohen-Macaulay rings,* Revised Edition, Cambridge, 1997.

[2] W. Bruns, R. Koch, *Normaliz*–a program for computing normalizations of affine semigroups, 1998. Available via anonymous ftp from: ftp.mathematik.Uni-Osnabrueck.DE/pub/osm/kommalg/software.

[3] J. Herzog, T. Hibi, Discrete polymatroids, *J. Algebraic Combin.*, **16**(2002), no. 3, 239–268.

[4] J. Herzog, T. Hibi, M. Vladoiu, Ideals of fiber type and polymatroids, Osaka J. Math. 42 (2005), 807-829.

[5] A. Ştefan, *A class of transversal polymatroids with Gorenstein base ring,* Bull. Math. Soc. Sci. Math. Roumanie Tome 51(99) No. 1, 2008.

[6] R. Villarreal, *Monomial Algebras,* Marcel Dekker, New-York, 2001.

[7] M. Vladoiu, Discrete polymatroids, An. Şt. Univ. Ovidius, Constanţa, 14 (2006), 89-112.

[8] M. Vladoiu, Equidimensional and unmixed ideals of Veronese type, to appear in Communications in Alg., arXiv:math. AC/0611326.

Alin Ştefan, Assistant Professor
"Petroleum and Gas" University of Ploieşti
Ploieşti, Romania
E-mail:nastefan@upg-ploiesti.ro